

The number counts, luminosity functions and evolution of microwave-selected (WMAP) blazars and radio galaxies

Paolo Giommi^{1,2}, Sergio Colafrancesco^{1,2,3}, Paolo Padovani⁴, Dario Gasparrini^{1,5,6},
Elisabetta Cavazzuti^{1,2}, Sara Cutini^{1,5,6}

¹ ASI Science Data Center, ASDC c/o ESRIN, Via G. Galilei 00044 Frascati, Italy.

² Agenzia Spaziale Italiana, Unità Osservazione dell'Universo, Viale Liegi 26 00198 Roma, Italy

³ on leave from: INAF - Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monteporzio, Italy

⁴ European Organisation for Astronomical Research in the Southern Hemisphere (ESO), Karl-Schwarzschild-Str. 2, D-85748 Garching bei München, Germany

⁵ INAF personnell resident at ASDC under ASI contract I/024/05/1

⁶ Department of Physics, University of Perugia, Via A. Pascoli, I-06123, Perugia, Italy

Received ; Accepted

Abstract. We have carried out an extensive search to identify the counterparts of all the microwave foreground sources listed in the WMAP 3-year catalogue using literature and archival data. Our work has led to the identification of 309 WMAP sources, 98% of which are blazars, radio quasars or radio galaxies. Only 7 WMAP detections were identified with other types of cosmic sources (3 starburst galaxies and 4 planetary/LBN nebulae). At present, 15 objects (< 5%) still remain without identification due to the unavailability of optical spectroscopic data or a clear radio counterpart. Our results allow us to define a flux limited sample of 203 high Galactic latitude microwave sources ($f_{41GHz} \geq 1$ Jy, $|b_{II}| > 15^\circ$) which is virtually completely identified (99%). The microwave band is ideally suited for blazar statistical studies since this is the part of the electromagnetic spectrum that is least affected by the superposition of spectral components of different origin, and therefore by selection effects. Using this data-set we have derived number counts, luminosity functions and cosmological evolution of blazars and radio galaxies at microwave frequencies. Our results are in good agreement with those found at radio (cm) frequencies. The 5 GHz bivariate blazar luminosity functions are similar to those derived from the DXRBS survey, which shows that this sample is representative of the blazar population at 41 GHz. Microwave selected broad-lined quasars are about six times more abundant than BL Lacs, a ratio that is similar to, or larger than, that seen at radio and gamma-ray frequencies, once spectral selection effects are taken into account. This strongly suggests that the mechanism responsible for the generation of gamma-rays is, at first order, the same in all blazar types, leaving little room for models (like external Compton radiation) that predict very different gamma-ray emission in broad-lined and lineless blazars. Our results confirm, and strengthen on a more solid statistical base, the findings of Giommi & Colafrancesco (2004, 2006) that blazars and radio galaxies are the largest contaminants of the CMB anisotropy maps. We predict that these sources are also bright gamma-ray sources, most of which will be detected by the AGILE and *Fermi* satellites.

Key words. galaxies: active – galaxies: Blazar: BL Lacertae surveys: Radio galaxies

microwave, hard X-ray, gamma-ray, TeV) are becoming available for deep astronomical observations.

1. Introduction

Blazars are the rarest and most peculiar type of Active Galactic Nuclei (AGN). So far almost 3,000 such objects have been reported in the literature (see e.g., Massaro et al. 2009 for an updated catalogue) but their number is bound to increase significantly in the near future as new, previously almost unexplored energy windows (e.g.

Blazar observational properties typically include irregular, sometimes large, rapid variability, apparent superluminal motion, flat radio spectrum, large and variable polarization at radio and, especially, optical frequencies. Because of their special observational properties, blazars are assumed to be sources emitting a continuum of electromagnetic radiation from a relativistic jet that is viewed closely along the line of sight thus causing strong relativistic amplification (e.g., Blandford & Rees 1978, Urry & Padovani 1995).

Blazars are a small fraction of all extragalactic sources but, unlike most other objects, they are strong emitters across the entire electromagnetic spectrum. In the optical and soft X-ray bands – where the radiation that we observe is mostly due to thermal emission originated in stars and galaxies or from the accretion process onto the central engine of AGN – blazars are a tiny minority, but in other parts of the electromagnetic spectrum, where thermal emission becomes unimportant, they are often the dominant population in the extragalactic sky. Following the technological evolution and the availability of astronomical resources, most blazars have so far been discovered as counterparts of flat spectrum radio emitters or as X-ray sources. Flux limited samples have been compiled from several surveys in both bands (e.g., Padovani et al. 2007 and references therein). In both these spectral regions, however, there are complications that may lead to selection effects of different nature: contamination by extended radio emission from the radio-lobes at cm wavelengths and contamination by radiation produced in the accretion process in the soft X-rays, among others. This may be particularly important in medium-low luminosity Flat Spectrum Radio Quasars (FSRQs) where the thermal emitted power from accretion processes and the non-thermal emission processes may be of comparable importance (e.g. Landt et al. 2008). The level of both thermal and non-thermal components can be in fact comparable also in the case of bright objects like, e.g., 3C 273 (see Grandi & Palumbo 2004).

The spectral energy distribution (SED) of blazars includes a synchrotron low-frequency component that peaks (in a $\text{Log}(\nu f(\nu)) - \text{Log}(\nu)$ representation) between the far infrared and the X-ray band, followed by an Inverse Compton high-frequency component that has its maximum in the hard X-ray band or at higher energies (see, e.g. Giommi et al. 2007), depending on the location of the synchrotron peak, and extends into the γ -ray or even in the TeV band.

Only recently the microwave region of the electromagnetic spectrum has become available to allow for systematic studies of blazars over large cosmological volumes. This frequency band is indeed particularly suited for the selection of blazars since at these frequencies the contamination from radio extended components with steep spectra is no longer present and the emission from the accretion process is negligible.

In this paper we present the first flux-limited sample of microwave selected blazars extracted from the catalogue of bright foreground sources detected by the WMAP satellite (Bennett et al. 2003, Hinshaw et al. 2007). Using this statistically complete sample we derive the number counts, the luminosity functions and the cosmological evolution of blazars and radio galaxies in the microwave band and we compare these properties with those observed at cm frequencies.

Throughout this paper, source spectra are written as $S_\nu \propto \nu^{-\alpha}$ where α is the spectral index. We use a flat, vacuum-dominated CDM cosmological model with

$H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$ (Spergel et al. 2006). To compare some of our results with previous works, we will also adopt a value $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and an empty universe cosmology with $\Omega_M = 0$, and $\Omega_\Lambda = 0$, as specifically declared in the text.

2. The sample

The catalogue of WMAP bright foreground sources that we consider here is based on the WMAP 3-year data (Hinshaw et al. 2007) and includes 324 sources detected by WMAP in at least one channel after three years of operation. Because of the higher sensitivity of the 3-year maps this catalogue is a significant improvement compared to that based on the first year data (Bennett et al. 2003) that included 208 sources.

For the purpose of this paper we will define our flux limited sample using the source fluxes as observed in the WMAP 41 GHz channel as a compromise between sensitivity and completeness, and the need to use the highest frequency band for the definition of a complete sample suitable for statistical purposes.

To arrive at the definition of such a complete sample, we first discuss the effect of source confusion in the estimate of the 41 GHz flux of the objects likely associated to the WMAP foreground sources. Then, we provide a classification scheme for each object associated to the WMAP source and finally we derive the statistical properties of our complete sample.

2.1. Source confusion

The WMAP experiment has a limited angular resolution which ranges between ~ 0.93 deg to 0.23 deg (FWHM) for the frequency bands 23 GHz to 94 GHz¹. In particular, the 41 GHz WMAP channel, to which we refer in our study, has an angular resolution of 31.8 arcmin (FWHM). Although this limited resolution, combined with the surface density of radio sources at around 1 Jy , does not imply a large fraction of confused sources, in some cases the probability that more than one object (especially those with flat radio spectra, like blazars) falls within the WMAP 41 GHz beam may not be negligible. When this happens the microwave flux attributed to a WMAP foreground source may be contaminated. We analyzed all the WMAP beams centered on the point-like sources detected at 41 GHz in order to find the level of confusion produced by the presence of multiple radio sources in the same beam, taking into consideration the effect of primary beam attenuation (Page et al. 2003). To this purpose, we adopted the following procedure:

- we started our analysis by associating the WMAP foreground source to the radio source listed in the WMAP 3 year catalogue (Hinshaw et al. 2007). However, in some cases we found arguments to propose a differ-

¹ <http://map.gsfc.nasa.gov/m-mm/ob-techres.html>

ent counterpart, as specifically discussed in Sect. 2.3 below;

- we choose to select radio sources potentially contaminating the 41 GHz flux as all sources within the WMAP 41 GHz beam centered on the object associated to the WMAP source that have radio flux at 5 GHz, $f_{5\text{GHz}} > 100$ mJy;
- for each radio source detected at 5 GHz within the WMAP beam, we estimated its 41 GHz flux by extrapolating the observed flux at lower frequencies using a linear regression (i.e., a power-law spectrum) technique. To this aim we used a variety of (non-simultaneous) radio catalogues including SUMSS, NVSS, GB6, PMN, and CRATES. Finally, we corrected the central WMAP source flux by subtracting the 41 GHz flux contributions of the possible contaminating objects taking into account the distance-dependent primary beam attenuation of the WMAP 41 GHz channel.

As the minimum error on the WMAP flux is 0.1 Jy, the correction was applied only when larger than this value. Our method could suffer from variability of the contaminants, which are mostly flat-spectrum radio sources, but this is unlikely to be a major problem because of the small fraction of “corrected” sources and amount of contamination. The percentage of sources, which ended up having their 41 GHz flux corrected, in fact, is only $\sim 7\%$, with the mean correction being $\sim 13\%$.

An example of a WMAP source which results confused by another flat-spectrum radio source in the field is shown in Fig. 1. This field includes two flat spectrum radio sources in the WMAP 41-GHz channel beam (the red circle with radius FWHM = 31.8 arcmin): the central source is located 1.8 arcsecs from the center of the field and has a flat spectrum with slope $\alpha_{1.4-4.8\text{GHz}} = -0.05$; there is a second bright flat spectrum source at 15 arcmin from the center of the field.

Fig. 2 shows a typical case in which there is no doubt about the association between the WMAP source and its radio counterpart. In this case, there is only one flat-spectrum ($\alpha_{1.4-4.8\text{GHz}} = 0.1$) radio source in the WMAP 41-GHz channel beam and it is very close (1.8 arcsecs) to the center of the field.

2.2. Object classification

We follow standard criteria and classify our sources based on their optical and radio spectra. The blazar class includes BL Lacertae objects, historically characterized by an almost complete lack of emission lines, and FSRQs, which by definition display broad, strong emission lines. A dividing value of full-width half maximum $FWHM = 1000$ km/s between “narrow” and “broad” emission lines is typically used. BL Lacs and radio galaxies can both have narrow lines so their separation is typically done on the basis of the value of the Ca break, a stellar absorption fea-




Fig. 1. An example of source confusion in the field of WMAP3 J0339-0144. Light grey filled circles represent radio sources (NVSS) either with steep radio spectrum or without spectral information; flat-spectrum radio sources are shown as dark blue filled circles. The green open circle marks the galaxy cluster ZW0334-0237 which is located 52 arcmin away from the center of the field. The size of the gray and blue circles is proportional to 1.4 GHz radio flux of the source they represent.




Fig. 2. Same as Fig. 1 but for the field of WMAP3 J2348-1631. The various sources in the field are NVSS sources either with steep radio spectrum or without spectral information (gray open circles) and flat-spectrum radio sources (blue open circles). The size of the gray and blue circles is proportional to 1.4 GHz radio flux. The WMAP source is clearly identified with the bright flat spectrum radio source (the blazar PKS 2345-16) at the center of the field.

ture in the optical spectrum defined by $C = (f_+ - f_-)/f_+$ (where f_- and f_+ are the fluxes in the rest frame wavelength regions 3750 – 3950 Å and 4050 – 4250 Å respectively). A separation value of $C = 0.4$ is normally adopted

(e.g., Padovani et al. 2007), as sources with $C < 0.4$ are believed to be dominated by non-stellar emission. We also make the commonly accepted distinction between steep spectrum radio quasars (SSRQ), $\alpha_r > 0.5$, and FSRQ, $\alpha_r \leq 0.5$. In order to separate BL Lacs (with broad emission lines) from radio quasars a limit of $5 \text{ \AA}$ on the rest frame equivalent width of any detected emission line is normally used (e.g., Stickel et al. 1991).

As most of the objects in the WMAP 3-year catalogue are bright and well known radio sources, the literature is rich with data. We have used the broad-band SED as well as any optical spectroscopy information available in the literature (mostly taken from the recent blazar catalog of Massaro et al. 2009, from the NASA Extragalactic Database and from the SDSS-DR6 on-line services) to classify all sources in the WMAP 3-year catalogue.

In summary, a WMAP source is associated to:

- an FSRQ if its spectral index at cm frequencies α_r and between the radio and the microwave region $\alpha_{1.4-41\text{GHz}}$ is flatter than 0.5, and if its optical spectrum shows strong and broad emission lines typical of QSOs, as detailed above;
- a BL Lac if its spectral index at cm frequencies α_r and between the radio and the microwave region $\alpha_{1.4-41\text{GHz}}$ is flatter than 0.5 and if its optical spectrum does not show strong broad emission lines and looks non-thermal, as detailed above;
- an SSRQ if its spectral index at cm frequencies α_r is steeper than 0.5 (independently of the $\alpha_{1.4-41\text{GHz}}$ index, which could be significantly flatter than α_r) and if its optical spectrum shows strong and broad emission lines;
- a radio galaxy if its spectral index at cm frequencies α_r is steeper than 0.5 (independently of the $\alpha_{1.4-41\text{GHz}}$ index), its radio emission is clearly extended, it has weak, narrow lines and an optical spectrum dominated by stellar emission, and if the host galaxy is clearly visible at optical/IR frequencies.

In a few cases we could not find sufficient information to classify a source with blazar characteristics as an object belonging to one of the above categories. In the following these sources are labelled as Blazars - Unknown type. Finally, in three cases the WMAP source appears to be classified in the literature as a starburst galaxy.

High frequency surveys detect a population of GigaHertz Peaked Spectrum (GPS) sources (e.g., O’Dea 1998). We have cross-correlated the WMAP list with the recent compilation by Labiano et al. 2007 and found 11 GPS sources, 5 of which belonging to the complete sample. These objects are marked accordingly in Tab. 3.

2.3. Notes on individual objects

WMAP3 J0026-3511 (complete sample). There are

no optical spectra of this object available in the literature. Its very flat radio spectrum and its blazar-like SED make this source a high-confidence blazar candidate. This source is very weak at low frequencies with a flux of only 14 mJy at 843 MHz (from SUMSS) and 25 mJy at 1.4 GHz (from NVSS). It is listed in the CRATES catalogue (Healey et al. 2007) as a very inverted radio source (spectral slope $\alpha_r = -1.4$) with a flux of 121 and 314 mJy at 4.8 and 8.4 GHz, respectively.

WMAP3 J0210-5100 (complete sample). This source is a transition object between FSRQs and BL Lac (Massaro et al. 2009). Given the relatively high redshift of this object ($z = 1$), we include it in the FSRQ sample.

WMAP3 J0540-5416, J0550-5732, J0633-2217, J1038+0511, J1102-4400, J1333+2723, J2035-6845 and J2148-7757 (complete sample: 1st, 2nd, and 4th source). These are all FSRQs that have been very recently identified by Healey et al. 2008 as part of the identification program of the CGRaBS gamma-ray source candidates.

WMAP3 J1149-7932. The object associated to this source in the WMAP 3 year catalog is not the most likely counterpart since it is too faint and with a rather steep spectrum.

WMAP3 J1227+1124. The position of the WMAP source does not correspond to a definite radio source. The GB6 source in the field is rather faint (34 mJy) and with flat spectrum. There is, however, a bright radio source associated to an interactive radio-galaxy (VPCX 27) at ≈ 25 arcmin from the nominal position of the WMAP 3 year source.

WMAP3 J1231+1351. Hinshaw et al. (2007) associate this source to GB6 J1231+1344 which is located 10 arcmin away from the WMAP source centroid. However, this radio source has a steep radio spectrum, while the source GB6 J1232+1359, at about 20 arcminutes from the WMAP centroid, is flat and brighter than GB6 J1231+1344. For this reason we associate WMAP3 J1231+1351 to GB6 J1232+1359.

WMAP3 J1305-4928. The WMAP source is associated to the galaxy NGC 4945, one of the four brightest FIR sources in the sky outside the local group. NGC 4945 hosts both vigorous nuclear star formation (Moorwood & Oliva 1994, Spoon et al. 2000) and a peculiar broad-lined AGN with an estimated bolometric luminosity of $\sim 60\%$ of the nuclear FIR luminosity. We classified this object as FIR-Starburst galaxy in our sample.

WMAP3 J1657+4749. We found 2 objects located inside the WMAP 41 GHz beam: 4C 48.41 and [HB89] 1656+477, both with a flat radio spectrum. The WMAP source has a 41 GHz flux of 0.6 Jy, but the two sources

mentioned above show a predicted flux at 41 GHz of 0.5 Jy and 1.3 Jy, respectively. Hinshaw et al. (2007) associate this source to the object GB6 1658+4737, that is [HB89] 1656+477. We agree with this preliminary association.

WMAP3 J1924-2914 (complete sample). Despite this source (PKS J1924-2914, PMN J1924-2914 or OV-236) being one of the brightest extragalactic objects at mm frequencies, it was not included in the WMAP 3-year catalog of Hinshaw et al. (2007) because it falls within the Kp0 cut that is used to avoid including a large number of Galactic sources in the catalog. Since the Kp0 cut is based on WMAP K-band intensity and this source is so bright, it falls within that cut even though it is a relatively high latitude source ($b_{\text{II}} = -19.6^\circ$; this is the only high-latitude extragalactic source to suffer this fate: G. Hinshaw private communication (see also Fig. 2 of Bennett et al. 2003). PKS J1924-2914, a well known, bright FSRQ, was clearly detected in the WMAP maps in all bands. Therefore, we have included it in our blazar sample. We estimated (from an analysis of the WMAP 3-yr data) that this source has a flux of $f_{41\text{GHz}} = 13.29 \pm 1.12$ Jy (with a mm spectral index $\alpha = 0.6$) and so it has been included in our complete sample.

WMAP3 J2333-2340 (complete sample). The optical spectrum of this source (Wilkes et al. 1983) shows emission lines and therefore, for consistency with current literature classification methods, we classify it as a radio galaxy. However the SED is typical of a blazar at all frequencies and shows a large optical variability. In a high optical state this object would have been classified as a BL Lac.

2.4. The complete sample

The full list of WMAP sources is given in Table 3 where column 1 gives the WMAP source name following the IAU naming convention, column 2 gives the WMAP number defined in the first year catalogue (if available), column 3 gives a common name (e.g. NGC, PKS, 3C etc.) if the source is a previously known object; columns 4 and 5 give the Right Ascension and Declination (J2000.0) of the counterpart of the WMAP source; column 6 gives the source class following the classification scheme described in Sect. 2.2; column 7 gives the redshift when available; columns 8 and 9 give the flux at 5 GHz (from the literature) and at 41 GHz (from the WMAP-3 yr catalogue); column 10 gives the 41 GHz flux corrected for source confusion as described in Sect. 2.1 and column 11 states if the source is part of the flux limited, complete sample.

The flux list of WMAP sources and the limited sample that we use for our statistical analysis is defined as the sample of all sources in the WMAP 3-yr catalogue with flux at 41 GHz larger than 1 Jy and with Galactic latitude $|b_{\text{II}}| > 15^\circ$. The latter condition is necessary to limit source confusion and the complications associated

to the emission close the Galactic plane. Taking also into account the higher latitude regions excluded by the Kp0 mask of Bennett et al. 2003 our total area is 28,457 square degrees. We chose $f_{41\text{GHz}} = 1$ Jy as the flux limit of our sample because the number of sources below this flux value drops sharply as shown in Fig. 3 where the distribution of 41 GHz fluxes is plotted for the entire sample. Above $F_{41\text{GHz}} = 1$ Jy (marked by the vertical dashed line) the slope of the distribution is steep and uniform indicating a very good level of completeness even close to the flux limit.

Table 1 gives the statistics of WMAP sources identification both for the full and the complete sample.

Source type	Number in sample total/complete
FSRQ	202/137
BL Lac	38/24
Blazars-unknown type	17/11
SSRQ	14/9
Radio galaxies	31/15
Starburst galaxies	3/2
Planetary/LBN nebulae	4/2
Unidentified/no radio-counterpart	15/3
Total	324/203

Table 1. Summary of WMAP objects identification

3. The blazar number counts at microwave frequencies (41 GHz)

In this section we present the 41 GHz logN-logS of blazars derived from our complete sample and we compare it to the counts derived at radio (5 GHz) frequency. Fig. 4 shows the integral LogN-LogS of *all* blazars (i.e. FSRQs, BL Lacs, and unclassified type) in the complete sample. The blazar counts at 41 GHz are steep and well described by a simple power law of the type $N(> S) \propto S^{-\beta}$. We use the maximum likelihood method (Crawford et al. 1970) to estimate the slope of the number counts using all the available information². This method yields $\beta = 1.64 \pm 0.14$, somewhat steeper than the Euclidean value of 1.5. The slopes for blazar sub-classes and other classes are consistent with this value within the typically large error bars.

This 41 GHz logN-logS is in very good agreement both in slope and normalization with the 5GHz blazar counts at flux densities > 1 Jy (see e.g. Giommi et al. 2006). This is expected given the flat $\langle \alpha_{5-41\text{GHz}} \rangle = 0.02$ average spectral slope of our blazars.

² This method operates on the differential distribution and obviates to the fact that individual points in integral counts are not independent.




Fig. 3. The distribution of the corrected 41 GHz flux in the entire WMAP sample. The sample appears to be clearly incomplete below 1 Jy, as marked by the dashed vertical line.




Fig. 4. The 41 GHz LogN-LogS of all blazars in the complete sample ($|b_{\text{II}}| > 15^\circ$) is shown by the filled dots with error bars. For clarity only a subset of the points is shown.

4. Cosmological evolution

The simplest way to study the evolutionary properties of a sample is through the V/V_{max} test (Schmidt 1968). Values of $\langle V/V_{\text{max}} \rangle$ significantly different from 0.5 indicate evolution, which is positive (i.e., sources were more luminous and/or more numerous in the past) for values > 0.5 , or negative (i.e., sources were less luminous and/or less numerous in the past) for values < 0.5 . One can also fit an evolutionary model to the sample by finding the evolutionary parameter which provides $\langle V/V_{\text{max}} \rangle = 0.5$.

We have computed the quantity V/V_{max} for our sources, with statistical errors given by $\sigma = 1/\sqrt{12 N}$ (Avni & Bahcall 1980). To have a simple estimate of the sample evolution we have also derived the best fit parameter τ (given in units of the age of the universe) of a pure lu-

minosity evolution model, i.e., $P(z) = P(0) \exp[T(z)/\tau]$, where $T(z)$ is the a-dimensional look-back time: the smaller τ is the stronger the evolution. We assume here that some luminosity evolution takes place, based on previous studies in the radio and other bands (e.g., Padovani et al. 2007, Croom et al. 2004).

The redshift distribution of WMAP blazars, SSRQs, radio galaxies is shown in Fig. 5.

The results of the cosmological evolution for our sample are shown in Table 2, which lists the source sub-sample in column (1), the number of sources in column (2), the mean redshift in column (3), the value of $\langle V/V_{\text{max}} \rangle$ and τ in columns (4) and (5) for our Λ CDM reference cosmological model, and the value of $\langle V/V_{\text{max}} \rangle$ and τ in columns (6) and (7) for an empty Universe cosmology, for comparison with previous results.




Fig. 5. The redshift distributions of blazars (FSRQs and BL Lacs), SSRQs, and radio galaxies.

The main results of our analysis can be summarized as follows:

1. FSRQs evolve at the 4.6σ level ($\langle V/V_{\text{max}} \rangle = 0.614 \pm 0.025$), confirming a well known result (e.g., Padovani & Urry 1992, Urry & Padovani 1995, Padovani et al. 2007). The evolutionary parameter of this class is consistent with that derived by Urry & Padovani 1995 for the 2 Jy FSRQ sample in an empty Universe ($\tau = 0.23^{+0.07}_{-0.04}$), for the simple case of pure luminosity evolution.
2. BL Lacs display moderate evolution, i.e., their value of $\langle V/V_{\text{max}} \rangle = 0.62 \pm 0.06$ is different from 0.5 at the 2σ level. The fact that 25% of the sources have no redshift information is not much of a problem, as redshift affects V/V_{max} values way less than flux.

3. Non blazars are consistent with no evolution (radio galaxies $\langle V/V_{\max} \rangle = 0.61 \pm 0.07$ and SSRQ $\langle V/V_{\max} \rangle = 0.63 \pm 0.10$). However, this is likely due to small number statistics, as SSRQ and high-power radio galaxies are known to evolve strongly in the radio band (e.g., Urry & Padovani 1995).
4. Blazars of unknown type show hints of evolution ($\langle V/V_{\max} \rangle = 0.70 \pm 0.09$) at the 2.3σ (2.5σ for an empty Universe cosmology) level, with a best fit value of τ consistent, within $\sim 1.5\sigma$, with that of FSRQs.

Given the available FSRQ statistics, we can move beyond the assumption of pure luminosity evolution and study possible redshift dependencies. We do this by using the so-called banded $\langle V/V_{\max} \rangle$ statistic, i.e. the quantity $\langle (V - V_o)/(V_{\max} - V_o) \rangle$, where V_o is the cosmological volume enclosed by a given redshift z_o (Dunlop & Peacock 1990). This procedure allows the detection of any high-redshift (possibly negative) evolution by separating it by the well-known strong (positive) low-redshift evolution. Fig. 6 shows that $\langle (V - V_o)/(V_{\max} - V_o) \rangle$ is basically constant up to $z \sim 1.5$ with a significant drop at higher redshifts. Its value, in fact, goes from being > 0.5 at the 4.6σ level at $z \sim 0$ to values ≤ 0.5 for $z \gtrsim 2$. However, given that only $\sim 10\%$ of the FSRQ are above this redshift, we assume a pure luminosity evolution for the whole sample. This is confirmed, within the somewhat limited statistics, by a study of the LF in redshift bins.




Fig. 6. The banded statistic, $\langle (V - V_o)/(V_{\max} - V_o) \rangle$ versus z_o for WMAP FSRQ. The horizontal dashed line indicates the value of 0.5 expected under the null hypothesis of no evolution.

5. The luminosity function of WMAP blazars

5.1. BL Lacs

The local LF of WMAP BL Lacs is shown in Fig. 7. Based on the $\langle V/V_{\max} \rangle$ analysis, this has been de-evolved to zero redshift using $\tau = 0.31$. We have assumed $z = \langle z \rangle \sim 0.56$ for the 6/24 BL Lacs without redshift.

We fitted the LF with a single power law $\phi(P_r) \propto P_r^{-B_r}$. Varying the luminosity binning, the differential slope is in the range $2.4 < B_r < 2.7$. For a representative bin size of $\Delta \log P_r = 0.4$, a weighted least-squares fit yields $\phi(P_r) \propto P_r^{-2.62 \pm 0.19}$ ($\chi^2_\nu \sim 0.94$ for 5 degrees of freedom). The total number density of BL Lacs in the luminosity range $7 \times 10^{24} - 10^{28}$ W/Hz, derived independently of bin size from the integral LF, equal to $\sum 1/V_{\max}$, is 44 ± 29 Gpc $^{-3}$ (see eqs. 9 and 10 of Condon et al. (2002)).

We note our assumption about the missing redshifts could somewhat bias our LF. Therefore, we have checked this by assuming $z = 1$ for these sources. As discussed in Padovani et al. 2007, their featureless continuum might suggest a relatively high redshift. However, this assumption results in a LF, which is consistent with the previous one well within the errors, with $\phi(P_r) \propto P_r^{-2.51 \pm 0.19}$ ($\chi^2_\nu \sim 0.69$ for 5 degrees of freedom).




Fig. 7. The local, de-evolved 41 GHz luminosity function of BL Lacs (filled points with error bars). Error bars correspond to 1σ Poisson errors (Gehrels 1986). The solid line is a weighted least-squares fit to the data of the form $\phi(P_r) \propto P_r^{-2.62}$ (see text for details).

We have also derived the bivariate LF of WMAP BL Lacs at 5 GHz for an empty Universe cosmology, to compare it with previous determinations and with the predictions of unified schemes. This is presented in Fig. 8 (filled points), which shows also the DXRBS

(open triangles, Padovani et al. 2007), 1 Jy LF (open squares, Stickel et al. 1991) and the predictions of a beaming model based on the 1 Jy LFs and evolution (solid line, Urry & Padovani 1995). These predictions assume that Fanaroff-Riley type I (FR I; Fanaroff & Riley 1974) galaxies are the counterparts of BL Lacs with their jets on the plane of the sky and show what one should expect to find when reaching powers lower than those used to constrain the LF at the high end. For consistency with Stickel et al. 1991 we have de-evolved the WMAP BL Lacs assuming $\tau = 0.32$, which is within 0.5σ from our own value (see Tab. 2). We note that 14/34 of the 1 Jy BL Lacs belong also to the WMAP sample.




Fig. 8. The bivariate 5 GHz luminosity function of WMAP BL Lacs (filled points) and DXRBS (open triangles) compared to the predictions of a beaming model based on the 1 Jy luminosity function and evolution (solid line Urry & Padovani 1995). The open squares represent the 1 Jy luminosity function Stickel et al. 1991. Error bars correspond to 1σ Poisson errors Gehrels 1986. The DXRBS points have been corrected to take into account their somewhat different definition of a BL Lac (Padovani et al. 2007). For consistency with Stickel et al. 1991 and Urry & Padovani 1995 the WMAP luminosity function has been de-evolved to zero redshift and an $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0$, and $\Omega_\Lambda = 0$ cosmology has been adopted.

The LF of WMAP BL Lacs is in very good agreement, in the region of overlap, with the DXRBS and 1 Jy LFs. In particular, the WMAP and 1 Jy LFs cover roughly the same power range, a result that is expected given the relatively similar flux limits. The WMAP LF is also in good agreement with the predictions of unified schemes. For this cosmology the total number density of BL Lacs

in the luminosity range $2 \times 10^{25} - 4 \times 10^{27} \text{ W/Hz}$, derived independently of bin size from the integral LF, is $17 \pm 11 \text{ Gpc}^{-3}$ to be compared with the value of 40 Gpc^{-3} in the range $6 \times 10^{24} - 3 \times 10^{27} \text{ W/Hz}$ for the 1 Jy LF.




Fig. 9. The local, de-evolved 41 GHz luminosity function of WMAP FSRQs. Error bars correspond to 1σ Poisson errors (Gehrels 1986).

5.2. FSRQs

The local LF of WMAP FSRQs is shown in Fig. 9. Based on the $\langle V/V_{\text{max}} \rangle$ analysis this has been de-evolved to zero redshift using $\tau = 0.35$.

We fitted the LF with a single power law $\phi(P_r) \propto P_r^{-B_r}$. Varying the binning, the differential slope is in the range $2.1 < B_r < 2.5$. For a bin size $\Delta \log P_r = 0.4$, which is representative, a weighted least-squares fit yields $\phi(P_r) \propto P_r^{-2.3 \pm 0.1}$ ($\chi^2_\nu \sim 5.1$ for 6 degrees of freedom). A single power law cannot then reproduce the observed LF (see Fig. 9). Note however that the first low-power bin includes a single source (3C 120) and the somewhat large gap between the first and the second bin. (We note that other authors have classified 3C 120 as a broad line radio galaxy).

The total number density of FSRQs in the luminosity range $6 \times 10^{24} - 10^{28} \text{ W/Hz}$, derived independently of bin size from the integral LF, is $30 \pm 28 \text{ Gpc}^{-3}$. Excluding the lowest luminosity source one gets $1.6 \pm 0.4 \text{ Gpc}^{-3}$ in the luminosity range $7 \times 10^{25} - 10^{28} \text{ W/Hz}$.

As already done for BL Lacs, we have also derived the local bivariate LF of FSRQs at 5 GHz for an empty Universe cosmology, to compare it with previous determinations and with the predictions of unified schemes. Based on the $\langle V/V_{\text{max}} \rangle$ analysis this has been de-evolved using




Fig. 10. The local, de-evolved bivariate 5 GHz radio luminosity function of WMAP FSRQs (filled points) and DXRBS with $z \leq 1$ (open triangles) compared to the predictions of a beaming model based on the 2 Jy luminosity function and evolution (solid line Urry & Padovani 1995). The open squares represent the 2 Jy luminosity function. Error bars represent the sum in quadrature of the 1σ Poisson errors (Gehrels 1986) and the variations of the number density associated with a 1σ change in the evolutionary parameter τ . For consistency with Urry & Padovani 1995 an empty cosmology with $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0$, and $\Omega_\Lambda = 0$ has been adopted. Both beaming model and 2 Jy LF have been converted from 2.7 GHz assuming $\alpha_r = 0$.

the value $\tau = 0.26$. The bivariate LF is shown in Fig. 10 (filled points), which also shows the DXRBS (open triangles, Padovani et al. 2007) and 1 Jy LFs (open squares, Stickel et al. 1991), and the predictions of a beaming model based on the 2 Jy LF and evolution (solid line, Urry & Padovani 1995). Given that the DXRBS evolutionary parameter is epoch dependent and that therefore we cannot simply de-evolve the global LF to zero redshift, we have restricted ourselves to sources having $z \leq 1$ (see Padovani et al. 2007 for details). We note that 36/52 of the 2 Jy and 8/129 of the DXRBS FSRQ belong to the WMAP sample.

The WMAP LF is in very good agreement, in the region of overlap, with the DXRBS and 2 Jy LFs. In particular, the WMAP and 2 Jy LFs cover roughly the same power range, as expected given the relatively similar flux limits. The WMAP LF is also in relatively good agreement with the predictions of unified schemes. Namely, most bins agree within $1-2\sigma$ with the beaming model, with the only exception of the two bins around 10^{27} W/Hz .

5.3. Radio galaxies and SSRQs

The LF of WMAP radio galaxies is shown in Fig. 11. Based on the $\langle V/V_{\text{max}} \rangle$ analysis no evolution has been assumed. We fitted the LF with a single power law $\phi(P_r) \propto P_r^{-B_r}$. Varying the binning, the differential slope is in the range $2.4 < B_r < 2.7$. For a representative bin size $\Delta \log P_r = 0.5$, a weighted least-squares fit yields $\phi(P_r) \propto P_r^{-2.6 \pm 0.1}$ ($\chi^2_\nu \sim 0.17$ for 5 degrees of freedom). The total number density of radio galaxies in the luminosity range $2 \times 10^{23} - 10^{27} \text{ W/Hz}$, derived independently of bin size from the integral LF, is $10,500 \pm 5,400 \text{ Gpc}^{-3}$.

As regards SSRQs, their sample is too small to derive a meaningful LF. However, we can say that their total number density in the luminosity range $5 \times 10^{24} - 4 \times 10^{27} \text{ W/Hz}$, derived independently of bin size from the integral LF, is $91 \pm 59 \text{ Gpc}^{-3}$.




Fig. 11. The 41 GHz luminosity function of radio galaxies (filled points). Error bars correspond to 1σ Poisson errors (Gehrels 1986). The solid line is a weighted least-squares fit to the data of the form $\phi(P_r) \propto P_r^{-2.6}$ (see text for details).

6. Discussion and Conclusions

In this paper we have presented a detailed study of the counterparts of all the sources of the WMAP 3-year catalogue, which has led to the identification of 309 microwave-selected objects. We have found that only 7 sources (three planetary nebulae, one Lynds Bright Nebula and three starburst galaxies) are neither blazars nor non-thermal misdirected AGN (i.e. radio galaxies and steep-spectrum radio quasars). All the remaining 302 sources have been classified according to the scheme presented in Sect. 2.2. This result fully confirms the previous findings

of Giommi & Colafrancesco 2004 that the vast majority of WMAP detected sources are blazars or radio galaxies.

We have used the WMAP 3-year source catalogue, after correcting their fluxes for source confusion within the WMAP 41 GHz channel beam, to define a microwave flux-limited, virtually complete, sample of blazars with $F_{41\text{GHz}} > 1$ Jy. Only 3 out of the 203 objects in this sample still remain unidentified. This is the first statistically well defined and complete sample of blazars selected at microwave frequencies. The microwave frequency range is likely the best region of the electromagnetic spectrum to pursue statistical studies of blazars since it is least affected by the superposition of spectral components of different nature like e.g. steep radio emission from the extended part of the jet, non-nuclear optical emission from the host galaxy or optical/UV and soft X-ray emission from an accretion disk.

Our complete sample allows us to derive number counts, luminosity function and cosmological evolution of WMAP AGN. We found that the LogN-LogS of microwave-selected blazars is steep, $N(> S) \sim S^{-1.64}$, down to 1 Jy, consistently with that found at cm frequencies (Giommi & Colafrancesco 2004, 2005, Giommi et al. 2006).

The bivariate 5 GHz luminosity functions of FSRQs and BL Lacs are similar to those derived from the DXRBS (for FSRQs and BL Lacs Padovani et al. 2007) and from the 2-Jy and 1-Jy FSRQs and BL Lacs surveys, respectively. A beaming model (Urry & Padovani 1995) adapted to the 1-Jy and 2-Jy blazar surveys is able to reproduce the observed luminosity function of WMAP blazars down to $P_r(5\text{GHz}) \sim 10^{25}$ W/Hz.

The cosmological evolution of these two subclasses of AGN is consistent with the expectations based on radio/cm measurements, yielding evolution at the 2σ level for BL Lacs and a quite strong luminosity evolution with a time scale $\tau = 0.35^{+0.07}_{-0.05}$ ($\tau = 0.26^{+0.05}_{-0.03}$) for a best-fit Λ CDM universe (an empty universe) reference model.

The luminosity function of radio galaxies can be fit with a power law shape with slope in the range 2.4–2.7 in the luminosity interval $P_r(41\text{GHz}) \approx 10^{23.3} - 10^{27}$ W/Hz. As these sources reach the smallest powers, they have the largest number density. Therefore, they are bound to be the dominant non-thermal AGN in the microwave sky at low flux densities. We find here a value of $\langle V/V_{\text{max}} \rangle = 0.61 \pm 0.07$, consistent with no evolution. However, this is easily explained by a combination of small number statistics, as the sample includes only 15 sources, small redshift range, and a mix of sub-classes. A look at the radio morphology of our radio galaxies, from both the literature and the NASA Extragalactic Database (NED), shows in fact a prevalence of FR I sources (9) followed by FR IIs (4), one GigaHertz Peaked Spectrum (GPS) and one compact steep-spectrum object. While FR Is are thought not to be evolving, FR IIs appear to evolve almost as strongly as radio quasars (see Urry & Padovani 1995 and references therein). We find a hint of this in our

data as well, since $\langle V/V_{\text{max}} \rangle = 0.57 \pm 0.10$ for FR Is and $\langle V/V_{\text{max}} \rangle = 0.67 \pm 0.12$ for non-FR Is.

The present complete sample of microwave-selected non-thermal AGN will allow us to predict the blazar contribution to the CMB maps obtainable with coming microwave experiments like *Planck* (see e.g., the *Planck* Blue Book) and Olimpo (Masi et al. 2005). The results obtained from our WMAP sample suggest that FSRQs and BL Lacs will likely dominate the CMB anisotropy spectrum at high microwave fluxes, while the other non-thermal AGN with lower fluxes but higher space density will provide a relevant contribution to the CMB anisotropies at lower microwave fluxes, if their LogN-LogS remains steep, as suggested by previous analyses at radio frequency.

The large number of faint non-blazar AGN will further provide a substantial contribution to the diffuse light (effective temperature) of the CMB sky observable with up-coming all-sky surveys. The additional contribution of microwave faint starburst galaxies (which are only marginally present in the high-flux threshold WMAP sample) will likely provide a relevant additional contribution to the unresolved CMB foreground light. We will present a detailed analysis of the contribution of different classes of non-thermal AGN to the cosmic background radiation field in a forthcoming paper (Padovani et al. 2009, in preparation).

Detailed considerations about the contribution of the polarized non-thermal AGN to the CMB polarization anisotropy spectrum will also be presented elsewhere (Colafrancesco et al. 2009, in preparation).

All the non-thermal AGNs (blazars, SSRQs and radio galaxies) in our sample are bright, well known sources that have been detected in several energy bands (from radio to X-rays) and are also expected to be detected in the gamma-ray band by the AGILE and *Fermi* experiments (see below). In this respect, almost 50 WMAP sources have already a counterpart in the third EGRET catalogue of gamma-ray sources (Hartman et al. 1999). In particular, AGILE is expected to detect a similar number of blazars and will monitor their duty cycle.

Finally, we note that the ratio of the number of FSRQs and BL Lacs at radio or microwave frequencies is similar to that found at gamma-ray energies. In fact, this ratio is ≈ 6 in our sample (see Table 2), ≈ 8 in the 5GHz DXRBS survey Padovani et al. 2007, and $\approx 3 - 5$ for the EGRET blazars Sowards-Emmerd et al. 2004).

Since the radio fluxes of EGRET detected blazars are similar to those of sources detected in the WMAP survey, blazar count ratios can be in agreement only with scenarios where the production of gamma-rays (that is intimately related to radio/microwave photons in Synchrotron - Inverse Compton models) is symmetric with respect to the two blazar types or slightly favour BL Lacs. Mechanisms like Synchrotron-Inverse Compton with External components (see, e.g., Sikora & Madejski 2001 for a review), which strongly favour FSRQs cannot be therefore very common otherwise blazars of this type

would be much more abundant than BL Lacs at gamma-ray frequencies. The recent *Fermi* results appear to confirm this. In fact, Abdo et al. 2009 find an FSRQ/BL Lac ratio of 1.3. This relatively low ratio is due to a strong selection effect, which picks up radio-faint BL Lacs with the synchrotron peak at high energies characterized by a flat gamma-ray spectrum. By comparing FSRQs and BL Lacs with similar synchrotron peaks (Giommi et al. 2009), one obtains a ratio ≈ 4 , not too different from that of WMAP and EGRET. In any case, even *Fermi* sees no evidence of enhanced gamma-ray emission from FSRQs as compared to that from BL Lacs.

We can conclude that the microwave and gamma-ray energy bands are best suited to study non-thermal AGN as their emission at these frequencies is largely dominated by non-thermal radiation. The combined study of non-thermal AGN in the microwave and in the gamma-ray energy bands will probably offer a unique opportunity to understand many of the physical details of this class of BH-dominated cosmic structures.

Acknowledgements. We are grateful to F. Verrecchia for his technical support in the analysis of the WMAP source confusion. We acknowledge the use of data and software facilities from the ASI Science Data Center (ASDC), managed by the Italian Space Agency (ASI). This work benefits from ASI grant I/024/05/1. Part of this work is based on archival data and bibliographic information obtained from the NASA Extragalactic Database (NED) and from the Astrophysics Data System (ADS).

References

- Abdo, A. A. et al. 2009, *ApJ*, 700, 597
- Avni, Y. & Bahcall, J.N. 1980, *ApJ*, 235, 694
- Bennett, C. L. et al. 2003, *ApJS*, 148, 97
- Blandford, R.D. & Rees, M.J. 1978, in *Pittsburg Conference on BL Lac Objects*, Ed. A.M. Wolfe, Pittsburgh, University of Pittsburgh press, p. 328
- Crawford, D.F., Jauncey, D.L., & Murdoch, H.S. 1970, *ApJ*, 162, 405
- Colafrancesco, S. et al. 2009, in preparation
- Condon, J. J., Cotton, W. D., & Broderick, J. J. 2002, *AJ*, 124, 675
- Croom, S.M. et al. 2004, *MNRAS*, 349, 1397
- Dunlop, J.S. & Peacock, J.A. et al. 1990 *MNRAS*, 247, 19
- Fanaroff, B.L. & Riley, J.M. 1974, *MNRAS*, 167, 31
- Gehrels, N. 1986, *ApJ*, 303, 336
- Giommi, P. & Colafrancesco, S. 2004, *A&A*, 414, 7
- Giommi, P. & Colafrancesco, S. 2005, *Exp. Astron.*, 20, 31
- Giommi, P. et al. 2006, *A&A*, 445, 843
- Giommi, P. et al. 2007, *A&A*, 468, 571
- Giommi, P. et al. 2007, *A&A*, 468, 97
- Giommi, P. et al. 2009, *ApJ*, submitted
- Grandi, P. & Palumbo, G. 2004, *Sci.*, 306, 998
- Hartman, R.C. et al. 1990, *ApJS*, 123, 79
- Healey, S.E. et al. 2007, *ApJS*, 171, 61
- Healey, S.E. et al. 2008, *ApJS*, 175, 97
- Hinshaw, G. et al. 2007, *ApJS*, 170, 288
- Labiano, A., Barthel, P.D., O'Dea, C.P., de Vries, W.H., Pérez, I., & Baum, S.A. 2007, *A&A*, 463, 97
- Landt, H. et. al. 2008, *ApJ*, 676, 87
- Marchã, M. J. M., Browne, I. W. A., Impey, C. D. & Smith, P. S. 1996, *MNRAS*, 281, 425
- Massaro, E. et al. 2009, *A&A*, 495, 691 (<http://www.asdc.asi.it/bzcat>)
- Masi, S. et al. 2005, in *17th ESA Symposium on European Rocket and Balloon Programmes and Related Research*, Sandefjord, Norway. Ed.: Barbara Warmbein. ESA SP-590, Noordwijk: ESA Publications Division, 2005, p. 581 - 586
- Moorwood, A.F. & Oliva, E. 1994, *ApJ*, 429, 602
- O'Dea, C. P. 1998, *PASP*, 110, 493
- Page, L. et al. 2003, *ApJS*, 148, 39
- Padovani, P. et al. 2007, *ApJ*, 662, 182
- Padovani, P. & Urry, M. 1992, *ApJ*, 387, 449
- Padovani, P. et al. 2009, in preparation
- Planck: The Scientific Programme (also known as the Blue Book). European Space Agency. ESA-SCI(2005)-1. Version 2.
- Schmidt, M. 1968, *ApJ*, 151, 393
- Sikora, M. & Madejski, G. 2001, in *AIP Conf. proc.* 558, *High Energy Gamma-Ray Astronomy*, F.A. Aharonian & H.J.Voelk Eds. (New York: AIP), p.275
- Sowards-Emmerd, D., Romani, R. W., Michelson, P. F., & Ulvestad, J. S. 2004, *ApJ*, 609, 564
- Spergel, D.N. et al. 2006, *ApJ*, 148, 175
- Spoon, H.W.W. et al. 2000, *A&A*, 357, 898
- Stickel, M. et al. 1991, *ApJ*, 374, 431
- Urry, M. & Padovani, P. 1995, *PASP*, 107, 803
- Wilkes, B. J., Wright, A. E., Jauncey, D. L. & Peterson, B. A. 1983, *PASA*, 5, 2

Table 2. WMAP sample Evolutionary Properties

Class	Number	$\langle z \rangle$	$H_0 = 70, \Omega_M = 0.3, \Omega_\Lambda = 0.7$		$H_0 = 50, \Omega_M = 0, \Omega_\Lambda = 0$	
			$\langle V/V_{\max} \rangle$	τ	$\langle V/V_{\max} \rangle$	τ
FSRQ	137	1.13 ± 0.05	0.614 ± 0.025	$0.35^{+0.07}_{-0.05}$	0.621 ± 0.025	$0.26^{+0.05}_{-0.03}$
BL Lacs	24	0.56 ± 0.09^a	0.619 ± 0.059^b	$0.31^{+0.21}_{-0.08}$	0.626 ± 0.059^b	$0.24^{+0.15}_{-0.05}$
Radio Galaxies	15	0.08 ± 0.04	0.610 ± 0.074	$\dots^c$	0.610 ± 0.074	$\dots^c$
Unclassified blazars	11	0.77 ± 0.16^d	0.703 ± 0.087^e	$0.22^{+0.11}_{-0.04}$	0.713 ± 0.087^e	$0.17^{+0.07}_{-0.03}$
SSRQ	9	0.50 ± 0.14	0.626 ± 0.096	$\dots^c$	0.632 ± 0.096	$\dots^c$

^aExcluding the 6 sources without redshift^bAssuming $z = \langle z \rangle$ for the sources without redshift^c $\langle V/V_{\max} \rangle$ not significantly different ($< 2 \sigma$) from 0.5: no evolution assumed^dExcluding the 6 sources without redshift^eAssuming z equal to the mean blazar redshift for the sources without redshift

Table 3: WMAP3 catalogue: basic data, identifications and inclusion in the flux limited sample

Source name IAU Convention	WMAP ID 1year catalogue	Common name	Ra(J2000.0)	Dec(J2000.0)	Class	Redshift	Radio flux density 5 GHz (mJy)	WMAP flux density 41 GHz (mJy)	Corrected 41 GHz (mJy)	In Complete Sample
WMAP3 J0006−0622	WMAP 60	PKS0003-066	00 06 13.8	−06 23 35.0	FSRQ	0.347	2463	2300	2300	yes
WMAP3 J0012−3953	WMAP 202	PKS0010-401	00 12 59.8	−39 54 24.9	BL Lac	0	921	1000	1000	yes
WMAP3 J0019+2019	—	PKS0017+200	00 19 37.7	+20 21 45.0	BL Lac	0	710	900	900	no
WMAP3 J0019+2602	—	4C25.01	00 19 39.1	+26 02 44.9	FSRQ	0.284	435	900	900	no
WMAP3 J0025−2604	—	PKS0023-26	00 25 49.0	−26 02 12.9	Radio Galaxy	0.322	3745	0	0	no
WMAP3 J0026−3511	—	PMNJ0026-3512	00 26 16.3	−35 12 48.9	Unidentified	0	121	1200	1200	yes
WMAP3 J0029+0555	—	PKS0027+056	00 29 45.7	+05 54 41.0	FSRQ	1.317	348	1200	1100	yes
WMAP3 J0042+5208	—	3C020	00 43 08.7	+52 03 33.9	Radio Galaxy	0.174	4124	0	0	no
WMAP3 J0047−2515	WMAP 62	NGC253	00 47 33.1	−25 17 17.0	Starburst galaxy	0.001	2433	1100	1100	yes
WMAP3 J0049−5740	WMAP 179	PKS0047-579	00 49 59.4	−57 38 26.9	FSRQ	1.797	1338	1000	1000	yes
WMAP3 J0051−0927	WMAP 77	PKS0048-09	00 50 41.2	−09 29 04.9	BL Lac	0.537	931	800	800	no
WMAP3 J0050−0647	—	PKS0048-71	00 51 08.2	−06 50 02.0	FSRQ	1.975	841	800	800	no
WMAP3 J0050−4223	—	PKS0048-427	00 51 09.4	−42 26 33.0	FSRQ	1.749	926	900	900	no
WMAP3 J0106−4035	WMAP 171	PKS0104-408	01 06 45.1	−40 34 19.9	FSRQ	0.584	2584	1800	1800	yes
WMAP3 J0108+0135	WMAP 81	PKS0106+01	01 08 37.6	+01 34 59.9	FSRQ	2.099	2078	2300	2300	yes
WMAP3 J0108+1320	WMAP 79	3C033	01 08 52.7	+13 19 13.0	Radio Galaxy	0.059	3691	1000	900	no
WMAP3 J0115−0126	—	PKS0112-017	01 15 17.1	−01 27 05.0	FSRQ	1.365	1540	800	800	no
WMAP3 J0116−1137	—	PKS0113-118	01 16 12.4	−11 36 15.0	Blazar Unknown-type	0.67	1488	1200	1200	yes
WMAP3 J0121+1150	—	PKS0119+11	01 21 41.5	+11 49 50.0	FSRQ	0.57	1126	1200	1200	yes
WMAP3 J0125−0010	WMAP 86	PKS0122-00	01 25 28.8	−00 05 56.0	FSRQ	1.076	1234	1300	1300	yes
WMAP3 J0132−1653	WMAP 97	PKS0130-17	01 32 43.3	−16 54 47.9	FSRQ	1.02	746	1900	1900	yes
WMAP3 J0137+4752	WMAP 80	S40133+47	01 36 58.4	+47 51 29.0	FSRQ	0.859	2016	4200	4200	no
WMAP3 J0137−2428	—	PKS0135-247	01 37 38.2	−24 30 54.0	FSRQ	0.837	956	1700	1700	yes
WMAP3 J0152+2208	—	PKS0149+21	01 52 17.9	+22 07 06.9	FSRQ	1.32	1073	1200	1200	yes
WMAP3 J0204+1513	WMAP 92	4C15.05	02 04 50.2	+15 14 11.0	FSRQ	0.405	3073	1500	1500	yes
WMAP3 J0204+3213	WMAP 85	1JY0202+319	02 05 04.9	+32 12 29.9	FSRQ	1.446	934	1200	1200	yes
WMAP3 J0210−5100	WMAP 158	PKS0208-512	02 10 46.0	−51 01 01.9	FSRQ	0.999	3198	2800	2800	yes
WMAP3 J0220+3558	—	1Jy0218+357	02 21 05.4	+35 56 13.9	Blazar Unknown-type	0.685	1480	1000	1000	yes
WMAP3 J0223+4304	WMAP 84	3C66B	02 23 11.3	+42 59 30.9	Radio Galaxy	0.021	1866	1400	1200	yes
WMAP3 J0222−3441	WMAP 137	PKS0220-349	02 22 56.4	−34 41 29.0	FSRQ	1.49	911	0	0	no
WMAP3 J0231+1318	—	4C13.14	02 31 45.7	+13 22 54.9	FSRQ	2.059	2608	1300	1300	yes
WMAP3 J0237+2848	WMAP 93	4C28.07	02 37 52.3	+28 48 08.9	FSRQ	1.213	2794	3500	3500	yes
WMAP3 J0238+1636	—	PKS0235+164	02 38 38.8	+16 36 59.0	BL Lac	0.94	1935	1400	1400	yes
WMAP3 J0253−5441	WMAP 155	PKS0252-549	02 53 29.1	−54 41 50.9	FSRQ	0.539	1193	2800	2800	yes
WMAP3 J0303−6212	WMAP 162	PKS0302-623	03 03 50.5	−62 11 25.0	Blazar Unknown-type	0	1862	1400	1400	yes
WMAP3 J0308+0404	WMAP 102	NGC1218	03 08 26.2	+04 06 39.9	Radio Galaxy	0.028	3773	1400	1400	yes
WMAP3 J0309−6103	WMAP 160	PKS0308-611	03 09 55.9	−60 58 38.9	Blazar Unknown-type	0	1103	900	900	no
WMAP3 J0312−7647	WMAP 174	PKS0312-77	03 11 55.1	−76 51 51.0	FSRQ	0.223	694	1100	1100	yes
WMAP3 J0319+4131	WMAP 94	NGC1275	03 19 48.1	+41 30 42.0	Radio Galaxy	0.017	46894	7000	7000	no
WMAP3 J0322−3711	WMAP 138	NGC1316	03 22 41.7	−37 12 38.9	Radio Galaxy	0.006	71133	10500	10500	yes
WMAP3 J0325+2224	—	GB6J0325+2223	03 25 36.7	+22 23 59.9	FSRQ	2.06	817	1100	1100	yes
WMAP3 J0329−2353	WMAP 123	PKS0327-241	03 29 54.0	−23 57 09.0	FSRQ	0.895	866	1200	1200	yes
WMAP3 J0334−4007	WMAP 146	PKS0332-403	03 34 13.6	−40 08 25.0	BL Lac	0	1331	1600	1600	yes
WMAP3 J0336−1258	—	PKS0334-131	03 36 34.9	−13 02 03.9	FSRQ	1.303	534	1200	1200	yes
WMAP3 J0339−0144	WMAP 106	PKS0336-019	03 39 30.8	−01 46 36.0	FSRQ	0.852	3014	2700	2500	yes
WMAP3 J0340−2121	—	PKS0338-214	03 40 35.5	−21 19 31.0	BL Lac	0.223	933	1200	1200	yes
WMAP3 J0348−2747	WMAP 129	PKS0346-279	03 48 38.0	−27 48 12.9	FSRQ	0.991	1444	800	800	no
WMAP3 J0358+1029	—	3C098	03 58 54.4	+10 26 03.0	Radio Galaxy	0.03	3132	0	0	no
WMAP3 J0403−3604	WMAP 136	PKS0402-362	04 03 53.7	−36 05 02.0	FSRQ	1.417	1851	4500	4500	yes
WMAP3 J0405−1305	WMAP 114	PKS0403-13	04 05 34.0	−13 08 12.9	FSRQ	0.57	2748	1800	1800	yes
WMAP3 J0411+7655	WMAP 82	4C76.03	04 10 45.5	+76 56 44.9	Radio Galaxy/GPS	0.598	5620	800	800	no
WMAP3 J0407−3824	WMAP 141	PKS0405-385	04 06 58.9	−38 26 27.9	Blazar Unknown-type	1.285	830	1000	1000	yes
WMAP3 J0408−7503	—	PKS0410-75	04 08 48.4	−75 07 18.9	Radio Galaxy	0.693	4569	600	600	no
WMAP3 J0423+0219	—	PKS0420+022	04 22 52.2	+02 19 27.0	FSRQ	0	1222	0	0	no
WMAP3 J0423−0120	WMAP 110	PKS0420-01	04 23 15.7	−01 20 33.0	FSRQ	0.914	4357	10100	10100	yes
WMAP3 J0424+0036	WMAP 109	PKS0422+004	04 24 46.7	+00 36 06.0	BL Lac	0.31	583	1800	1700	yes
WMAP3 J0424−3757	WMAP 140	PKS0422-380	04 24 42.1	−37 56 20.0	FSRQ	0.782	1706	1300	1300	yes

Giommi et al.: A WMAP complete sample of blazars

Table 3: WMAP3 catalogue: basic data, identifications and inclusion in the flux limited sample (continued)

Giommi et al.: A WMAP complete sample of blazars

Source name IAU Convention	WMAP ID 1year catalogue	Common name	Ra(J2000.0)	Dec(J2000.0)	Class	Redshift	Radio Flux density 5 GHz (mJy)	WMAP flux density 41 GHz (mJy)	Corrected 41 GHz (mJy)	In Complete Sample
WMAP3 J0428−3757	—	PKS0426-380	04 28 40.3	−37 56 20.0	BL Lac	1.11	1202	1300	1300	yes
WMAP3 J0433+0521	WMAP 108	3C120	04 33 10.9	+05 21 15.0	FSRQ	0.033	5189	2600	2600	yes
WMAP3 J0440−4333	WMAP 147	PKS0438-43	04 40 17.1	−43 33 09.0	FSRQ	2.863	3933	2800	2800	yes
WMAP3 J0442−0018	—	PKS0440-00	04 42 38.5	−00 17 43.0	FSRQ	0.844	1084	1200	1200	yes
WMAP3 J0453−2806	WMAP 131	PKS0451-28	04 53 14.5	−28 07 36.9	FSRQ	2.559	2172	0	0	no
WMAP3 J0449−8100	WMAP 175	PKS0454-81	04 50 05.3	−81 01 01.9	FSRQ	0.444	1357	1700	1700	yes
WMAP3 J0455−4617	WMAP 151	PKS0454-46	04 55 50.7	−46 15 59.0	FSRQ	0.853	1653	4000	4000	yes
WMAP3 J0457−2322	WMAP 128	PKS0454-234	04 57 03.1	−23 24 51.9	FSRQ	1.003	1863	2600	2600	yes
WMAP3 J0501−0158	—	4C-02.19	05 01 12.7	−01 59 13.9	FSRQ	2.286	3317	1300	1200	yes
WMAP3 J0506−6108	WMAP 154	PKS0506-61	05 06 43.9	−61 09 41.0	FSRQ	1.093	1211	1700	1500	yes
WMAP3 J0513−2155	WMAP 127	PKS0511-220	05 13 49.0	−21 59 16.0	FSRQ	1.296	865	1000	1000	yes
WMAP3 J0513−2015	—	PMNJ0514-2029	05 14 17.2	−20 29 21.0	Unidentified	0	431	800	700	no
WMAP3 J0515−4558	—	PKS0514-459	05 15 45.3	−45 46 44.0	FSRQ	0.194	0	900	400	no
WMAP3 J0519−0540	WMAP 116	—	05 19 14.5	−05 40 41.0	LBN Nebula	0	0	1800	1800	yes
WMAP3 J0519−4546	WMAP 150	PICTORA	05 19 49.6	−45 46 44.0	SSRQ	0.035	15827	4200	4200	yes
WMAP3 J0523−3627	WMAP 139	PKS0521-36	05 22 57.8	−36 27 30.9	BL Lac	0.055	8180	3400	3400	yes
WMAP3 J0525−4826	—	PKS0524-485	05 26 16.6	−48 30 35.9	Unidentified	0	425	1500	1500	yes
WMAP3 J0527−1240	WMAP 122	IC0418	05 27 28.2	−12 41 49.9	Planetary Nebula	0	1716	1400	1400	yes
WMAP3 J0538−4405	WMAP 148	PKS0537-441	05 38 50.2	−44 05 08.9	BL Lac	0.894	4805	5700	5700	yes
WMAP3 J0542+4951	WMAP 95	3C147.0	05 42 36.1	+49 51 06.9	SSRQ	0.545	7264	1100	1100	no
WMAP3 J0540−5416	WMAP 152	PKS0539-543	05 40 45.7	−54 18 21.9	FSRQ	1.185	373	1800	1800	yes
WMAP3 J0550−5732	WMAP 153	PKS0549-575	05 50 09.4	−57 32 23.9	FSRQ	2.001	355	1200	1200	yes
WMAP3 J0555+3944	WMAP 100	B20552+39A	05 55 30.7	+39 48 48.9	FSRQ	2.365	5524	1100	1100	no
WMAP3 J0600−4528	—	PKS0557-454	05 59 11.5	−45 29 39.9	FSRQ	0.687	407	900	900	no
WMAP3 J0607+6722	WMAP 91	S40602+673	06 07 52.0	+67 20 54.9	FSRQ	1.97	581	700	700	no
WMAP3 J0608−2219	—	PKS0606-223	06 08 59.7	−22 20 21.0	FSRQ	1.926	626	1000	1000	yes
WMAP3 J0609−1541	WMAP 126	PKS0607-15	06 09 40.9	−15 42 41.0	FSRQ	0.324	2473	3400	3400	yes
WMAP3 J0623−6436	—	IRASL06229-643	06 23 07.7	−64 36 20.9	FSRQ	0.129	531	900	900	no
WMAP3 J0629−1957	WMAP 130	PKS0627-199	06 29 23.7	−19 59 20.0	BL Lac	0	784	1600	1600	no
WMAP3 J0633−2217	WMAP 135	PMNJ0633-2223	06 33 26.7	−22 23 21.0	FSRQ	1.508	885	900	900	no
WMAP3 J0639+7326	WMAP 87	S50633+734	06 39 21.8	+73 24 57.9	FSRQ	1.85	711	1100	1100	yes
WMAP3 J0636−2032	WMAP 134	PKS0634-20	06 36 32.2	−20 34 53.0	Radio Galaxy	0.055	1326	900	900	no
WMAP3 J0635−7517	WMAP 167	PKS0637-75	06 35 46.4	−75 16 17.0	FSRQ	0.653	6398	4000	4000	yes
WMAP3 J0646+4449	WMAP 99	S40642+449	06 46 31.9	+44 51 15.9	FSRQ/GPS	3.396	1220	2200	2200	yes
WMAP3 J0721+7122	—	S50716+714	07 21 53.4	+71 20 35.9	BL Lac	0.3	859	2200	2200	yes
WMAP3 J0721+0401	—	GB6J0721+0406	07 21 23.8	+04 06 46.0	Unidentified	0	533	700	700	no
WMAP3 J0727+6748	—	3C179	07 28 11.2	+67 48 47.0	SSRQ	0.846	912	700	700	no
WMAP3 J0725−0049	—	PKS0723-008	07 25 50.5	−00 54 56.0	Blazar Unknown-type	0.128	1377	1200	1200	no
WMAP3 J0734+5021	—	GB6J0733+5022	07 33 52.4	+50 22 09.0	FSRQ	0.72	991	1200	1200	yes
WMAP3 J0738+1744	WMAP 113	PKS0735+17	07 38 07.3	+17 42 19.0	BL Lac	0.42	1812	1300	1200	yes
WMAP3 J0739+0136	WMAP 124	PKS0736+01	07 39 18.0	+01 37 04.0	FSRQ	0.19	1781	2500	2500	no
WMAP3 J0741+3111	WMAP 107	OI363	07 41 10.6	+31 11 59.9	FSRQ	0.63	2623	900	900	no
WMAP3 J0745+1012	WMAP 118	PKS0742+10	07 45 33.0	+10 11 12.9	FSRQ/GPS	2.624	3704	800	800	no
WMAP3 J0746−0045	—	4C-00.28	07 45 54.1	−00 44 17.0	FSRQ/GPS	0.994	1975	800	800	no
WMAP3 J0743−6727	WMAP 161	PKS0743-67	07 43 31.6	−67 26 25.0	FSRQ	1.51	2235	700	600	no
WMAP3 J0750+1230	WMAP 117	PKS0748+126	07 50 52.0	+12 31 05.0	FSRQ	0.889	1238	2600	2600	yes
WMAP3 J0753+5354	—	4C54.15	07 53 01.3	+53 52 59.0	BL Lac	0	964	1000	1000	yes
WMAP3 J0757+0957	WMAP 120	PKS0754+100	07 57 06.6	+09 56 35.0	BL Lac	0.266	1037	1600	1600	yes
WMAP3 J0808−0750	WMAP 133	PKS0805-07	08 08 15.4	−07 51 10.0	FSRQ	1.837	1599	1400	1400	no
WMAP3 J0813+4817	—	3C196	08 13 35.9	+48 13 00.9	SSRQ	0.871	4504	0	0	no
WMAP3 J0816−2425	WMAP 145	PMNJ0816-2421	08 16 40.3	−24 21 05.0	Unidentified	0	270	1200	1200	no
WMAP3 J0824+3914	—	4C+39.23	08 24 55.5	+39 16 41.9	FSRQ	1.216	1031	1000	900	no
WMAP3 J0825+0311	WMAP 125	PKS0823+033	08 25 50.2	+03 09 24.0	BL Lac	0.506	2064	1500	1500	yes
WMAP3 J0830+2410	WMAP 112	B20827+24	08 30 52.0	+24 10 59.9	FSRQ	0.94	886	1600	1600	yes
WMAP3 J0838+5820	—	S40833+58	08 37 22.3	+58 25 00.9	FSRQ	2.101	717	1000	1000	yes
WMAP3 J0836−2015	WMAP 144	PKS0834-20	08 36 39.1	−20 16 59.0	FSRQ	2.752	1712	2300	2300	no
WMAP3 J0841+7053	WMAP 89	4C71.07	08 41 24.2	+70 53 41.9	FSRQ	2.172	2342	1800	1800	yes
WMAP3 J0840+1312	WMAP 121	3C207.0	08 40 47.6	+13 12 23.0	SSRQ	0.681	1244	1800	1800	yes

Table 3: WMAP3 catalogue: basic data, identifications and inclusion in the flux limited sample (continued)

Source name IAU Convention	WMAP ID 1year catalogue	Common name	Ra(J2000.0)	Dec(J2000.0)	Class	Redshift	Radio Flux density 5 GHz (mJy)	WMAP flux density 41 GHz (mJy)	Corrected 41 GHz (mJy)	In Complete Sample
WMAP3 J0854+2005	WMAP 115	PKS0851+202	08 54 48.7	+20 06 29.9	BL Lac	0.306	2908	4200	4200	yes
WMAP3 J0902−1414	—	PKS0859-14	09 02 16.8	−14 15 30.9	FSRQ	1.333	2022	1300	1300	yes
WMAP3 J0907−2017	—	PMNJ0907-2026	09 07 54.1	−20 26 51.0	Unidentified	0	417	0	0	no
WMAP3 J0909+4252	—	3C216	09 09 33.4	+42 53 47.0	Radio Galaxy	0.67	1561	1200	1200	yes
WMAP3 J0909+0118	WMAP 132	4C01.24	09 09 10.0	+01 21 36.0	FSRQ	1.024	888	1800	1800	yes
WMAP3 J0914+0249	—	PKS0912+029	09 14 37.9	+02 45 59.0	FSRQ	0.427	842	1400	1400	yes
WMAP3 J0918−1203	WMAP 143	HYDRAA	09 18 05.6	−12 05 44.0	Radio Galaxy	0.055	13982	1000	1000	yes
WMAP3 J0921+6216	—	S40917+62	09 21 36.1	+62 15 51.9	FSRQ	1.446	1226	900	900	no
WMAP3 J0920+4440	—	S40917+44	09 20 58.3	+44 41 53.9	FSRQ	2.19	1085	1300	1300	yes
WMAP3 J0921−2619	—	PKS0919-260	09 21 29.4	−26 18 42.9	FSRQ	2.3	2333	1200	1200	yes
WMAP3 J0927+3901	WMAP 105	4C39.25	09 27 02.8	+39 02 21.0	FSRQ	0.695	6913	5200	5200	yes
WMAP3 J0948+4037	WMAP 104	4C40.24	09 48 55.3	+40 39 45.0	FSRQ	1.249	1801	1400	1400	yes
WMAP3 J0955+6936	WMAP 88	M82	09 55 51.7	+69 40 45.0	Starburst galaxy	0.001	3796	1100	1100	yes
WMAP3 J0957+5526	—	4C55.17	09 57 38.1	+55 22 58.0	FSRQ	0.896	2015	1000	1000	yes
WMAP3 J0958+4721	WMAP 98	S40955+476	09 58 19.6	+47 25 08.0	FSRQ	1.882	1005	1400	1400	yes
WMAP3 J1014+2304	WMAP 119	4C23.24	10 14 47.1	+23 01 15.9	FSRQ	0.565	1093	900	900	no
WMAP3 J1018−3129	—	PKS1016-311	10 18 28.6	−31 23 53.9	FSRQ	0.794	688	0	0	no
WMAP3 J1032+4118	WMAP 103	S41030+415	10 33 03.7	+41 16 05.9	FSRQ	1.117	439	900	900	no
WMAP3 J1037−2934	—	PKS1034-293	10 37 16.0	−29 34 02.9	Blazar Unknown-type	0.312	1477	1100	1100	yes
WMAP3 J1038+0511	WMAP 142	PKS1036+054	10 38 46.7	+05 12 29.0	FSRQ	0.473	704	1500	1500	yes
WMAP3 J1041+0611	—	4C06.41	10 41 17.0	+06 10 17.0	FSRQ	1.27	1518	1400	1400	yes
WMAP3 J1041−4740	WMAP 163	PMNJ1041-4740	10 41 44.2	−47 40 00.0	Unidentified	0	2480	300	300	no
WMAP3 J1047+7143	WMAP 83	S51044+719	10 48 27.6	+71 43 35.0	FSRQ	1.15	1900	800	800	no
WMAP3 J1047−1911	—	PKS1045-18	10 48 06.5	−19 09 36.0	FSRQ	0.595	879	800	800	no
WMAP3 J1058+0134	WMAP 149	4C01.28	10 58 29.5	+01 33 59.0	Blazar Unknown-type	0.89	3263	4500	4500	yes
WMAP3 J1059−8003	WMAP 176	PKS1057-79	10 58 43.3	−80 03 53.9	Blazar Unknown-type	0	2130	2200	2200	yes
WMAP3 J1102−4400	—	PKS1059-438	11 02 04.8	−44 04 22.0	FSRQ	0.763	448	0	0	no
WMAP3 J1107−4446	WMAP 166	PKS1104-445	11 07 08.5	−44 49 08.0	FSRQ	1.598	3325	1200	1200	no
WMAP3 J1117−4635	—	PKS1116-46	11 18 26.8	−46 34 14.9	FSRQ	0.713	1872	600	600	no
WMAP3 J1118+1238	—	4C12.39	11 18 57.3	+12 34 41.9	FSRQ	2.129	1820	1100	1100	yes
WMAP3 J1127−1858	WMAP 159	PKS1124-186	11 27 04.3	−18 57 16.9	FSRQ	1.05	1624	1300	1300	yes
WMAP3 J1130−1452	WMAP 157	PKS1127-145	11 30 07.0	−14 49 27.0	FSRQ/GPS	1.184	4209	1500	1500	yes
WMAP3 J1130+3813	WMAP 101	S41128+385	11 30 53.2	+38 15 18.0	FSRQ	1.741	769	1100	1100	yes
WMAP3 J1146−4838	—	PKS1143-48	11 45 30.7	−48 36 11.9	Radio Galaxy	0.33	1376	0	0	no
WMAP3 J1146+4001	—	S41144+402	11 46 58.2	+39 58 33.9	FSRQ	1.088	836	1100	1100	yes
WMAP3 J1147−3811	WMAP 169	PKS1144-379	11 47 01.3	−38 12 11.0	BL Lac	1.048	1825	1900	1700	yes
WMAP3 J1149−7932	—	PMNJ1147-7936	11 46 50.5	−79 37 09.0	Unidentified	0	0	700	700	no
WMAP3 J1154+8104	WMAP 78	S51150+81	11 53 12.4	+80 58 28.9	FSRQ	1.25	1180	0	0	no
WMAP3 J1153+4931	WMAP 90	4C49.22	11 53 24.4	+49 31 09.0	FSRQ	0.334	717	2200	1600	yes
WMAP3 J1159+2914	WMAP 111	4C29.45	11 59 31.7	+29 14 44.0	FSRQ	0.729	1461	2000	2000	yes
WMAP3 J1203+4806	—	GB6J1203+4803	12 03 29.8	+48 03 12.9	FSRQ	0.817	164	700	700	no
WMAP3 J1208−2405	WMAP 172	PKSB1206-238	12 09 02.3	−24 06 20.9	FSRQ	1.299	1073	900	900	no
WMAP3 J1215−1730	WMAP 173	PKS1213-17	12 15 46.7	−17 31 45.9	Blazar Unknown-type	0	1744	1200	1200	yes
WMAP3 J1218+4832	—	S41216+48	12 19 06.4	+48 29 56.0	FSRQ	1.076	638	800	800	no
WMAP3 J1219+0548	WMAP 164	NGC4261	12 19 23.2	+05 49 31.0	Radio Galaxy	0.007	9064	2000	2000	yes
WMAP3 J1222−8307	WMAP 178	PKS1221-82	12 24 54.4	−83 13 09.9	Blazar Unknown-type	0	797	900	900	no
WMAP3 J1227+1124	—	GB6J1228+1124	12 28 29.1	+11 24 52.9	Unidentified	0	0	500	500	no
WMAP3 J1229+0203	WMAP 170	3C273	12 29 06.7	+02 03 07.9	FSRQ	0.158	36923	15700	15700	yes
WMAP3 J1230+1223	WMAP 165	M87	12 30 49.3	+12 23 27.9	Radio Galaxy	0.004	59027	13100	13100	yes
WMAP3 J1231+1351	—	GB6J1231+1344	12 31 42.9	+13 44 48.0	No clear radio counterpart	0.115	0	0	0	no
WMAP3 J1246−2547	WMAP 177	PKS1244-255	12 46 46.6	−25 47 48.0	FSRQ	0.633	2317	1800	1800	yes
WMAP3 J1256−0547	WMAP 181	3C279	12 56 10.9	−05 47 21.0	FSRQ	0.536	11192	18000	18000	yes
WMAP3 J1258−3158	WMAP 180	PKS1255-316	12 57 58.9	−31 55 14.9	FSRQ	1.924	1410	1300	1300	yes
WMAP3 J1258−2223	—	PKS1256-220	12 58 54.4	−22 19 32.0	FSRQ	1.306	978	1000	1000	yes
WMAP3 J1302+4856	—	—	13 02 49.3	+48 56 35.9	No clear radio counterpart	0	0	900	900	no
WMAP3 J1305−4928	—	NGC4945	13 05 27.4	−49 28 04.0	FIR Starburst galaxy	0.002	3055	0	0	no
WMAP3 J1309+1155	—	MC21307+12	13 09 33.9	+11 54 24.9	BL Lac	0	1350	1000	1000	yes
WMAP3 J1310+3221	WMAP 52	1Jy1308+326	13 10 28.6	+32 20 43.0	BL Lac	0.996	1447	2700	2000	yes

Giommi et al.: A WMAP complete sample of blazars

Table 3: WMAP3 catalogue: basic data, identifications and inclusion in the flux limited sample (continued)

Source name IAU Convention	WMAP ID 1year catalogue	Common name	Ra(J2000.0)	Dec(J2000.0)	Class	Redshift	Radio Flux density 5 GHz (mJy)	WMAP flux density 41 GHz (mJy)	Corrected 41 GHz (mJy)	In Complete Sample
WMAP3 J1316−3337	WMAP 182	PKS1313-333	13 16 07.9	−33 38 58.9	FSRQ	1.21	1093	1700	1700	yes
WMAP3 J1324−1046	—	PKSB1321-105	13 24 25.7	−10 49 23.0	FSRQ	0.872	857	1000	1000	yes
WMAP3 J1327+2212	—	GB6J1327+2210	13 27 00.7	+22 10 50.0	FSRQ	1.4	647	800	800	no
WMAP3 J1329+3201	WMAP 40	GB6J1329+3154	13 29 52.7	+31 54 11.0	BL Lac	0	518	800	800	no
WMAP3 J1330+2504	—	3C287	13 30 37.6	+25 09 10.0	SSRQ	1.055	3289	1100	1100	yes
WMAP3 J1331+3031	WMAP 26	3C286	13 31 08.2	+30 30 32.0	SSRQ	0.849	6350	1400	1400	yes
WMAP3 J1332+0200	—	3C287.1	13 32 53.2	+02 00 45.0	SSRQ	0.216	1369	1100	1100	yes
WMAP3 J1333+2723	—	GB6J1333+2725	13 33 07.3	+27 25 18.0	FSRQ	2.126	318	900	900	no
WMAP3 J1336−3358	WMAP 185	IC4296	13 36 43.0	−33 58 40.0	Radio Galaxy	0.012	1886	1100	1100	yes
WMAP3 J1337−1257	WMAP 188	PKS1335-127	13 37 39.7	−12 57 24.0	FSRQ	0.539	2838	6400	6400	yes
WMAP3 J1343+6601	—	S41342+663	13 43 45.9	+66 02 26.0	FSRQ	0.776	545	600	100	no
WMAP3 J1354−1041	WMAP 197	PKS1352-104	13 54 46.9	−10 41 02.0	FSRQ	0.332	686	1600	1600	yes
WMAP3 J1357−1529	—	PKS1354-152	13 57 11.2	−15 27 29.0	FSRQ	1.89	841	1000	1000	yes
WMAP3 J1356+1919	WMAP 4	4C19.44	13 57 04.3	+19 19 06.9	FSRQ	0.72	2618	1200	1200	yes
WMAP3 J1357+7644	—	S51357+769	13 57 55.3	+76 43 19.9	Blazar Unknown-type	0	844	800	800	no
WMAP3 J1408−0749	WMAP 203	PKS1406-076	14 08 56.4	−07 52 27.0	FSRQ	1.494	1080	1100	1100	yes
WMAP3 J1411+5216	—	3C295	14 11 20.5	+52 12 09.0	Radio Galaxy	0.464	7374	0	0	no
WMAP3 J1419+3823	WMAP 42	B31417+385	14 19 46.6	+38 21 47.9	FSRQ	1.831	651	1000	1000	yes
WMAP3 J1420+2702	—	GB6J1419+2706	14 19 59.2	+27 06 24.9	FSRQ	0.536	415	1100	1000	yes
WMAP3 J1419+5425	—	1Jy1418+546	14 19 46.6	+54 23 13.9	BL Lac	0.153	1350	900	900	no
WMAP3 J1427−3303	WMAP 193	PKS1424-328	14 27 41.4	−33 05 30.9	Unidentified	0	352	1700	1600	yes
WMAP3 J1427−4206	WMAP 191	PKS1424-41	14 27 56.2	−42 06 19.0	FSRQ	1.522	2597	2900	2900	yes
WMAP3 J1440+4958	—	GB6J1439+4958	14 39 47.8	+49 58 05.0	BL Lac	0	198	0	0	no
WMAP3 J1442+5156	—	3C303	14 43 02.7	+52 01 36.9	SSRQ	0.141	1070	800	800	no
WMAP3 J1458+7140	WMAP 71	3C309.1	14 59 07.6	+71 40 18.9	SSRQ	0.905	3567	1100	1100	yes
WMAP3 J1504+1029	WMAP 6	PKS1502+106	15 04 24.9	+10 29 38.0	FSRQ	1.839	2325	1600	1600	yes
WMAP3 J1506−1644	—	PKS1504-167	15 07 04.8	−16 52 30.0	FSRQ	0.876	2840	1000	1000	yes
WMAP3 J1510−0548	—	4C-05.64	15 10 53.5	−05 43 06.9	FSRQ	1.185	1742	1400	1400	yes
WMAP3 J1512−0905	WMAP 207	PKS1510-08	15 12 50.5	−09 05 59.9	FSRQ	0.36	3080	1800	1800	yes
WMAP3 J1514−1013	—	PKS1511-10	15 13 44.8	−10 11 59.9	FSRQ	1.513	1220	1100	1100	yes
WMAP3 J1516+0013	WMAP 2	4C00.56	15 16 40.1	+00 15 02.0	Radio Galaxy	0.052	1641	1700	1700	yes
WMAP3 J1517−2421	WMAP 205	APLIB	15 17 41.8	−24 22 18.9	BL Lac	0.049	2013	1900	1900	yes
WMAP3 J1540+1446	—	4C14.60	15 40 49.5	+14 47 45.9	BL Lac	0.605	1210	800	800	no
WMAP3 J1549+0236	WMAP 5	PKS1546+027	15 49 29.4	+02 37 00.9	FSRQ	0.414	1078	2100	2100	yes
WMAP3 J1549+5037	—	S41547+507	15 49 17.5	+50 38 05.9	FSRQ	2.175	731	900	900	no
WMAP3 J1550+0526	WMAP 7	4C05.64	15 50 35.2	+05 27 10.0	FSRQ	1.422	1766	2000	2000	yes
WMAP3 J1556−7912	—	PKS1549-79	15 56 58.9	−79 14 03.9	Radio Galaxy	0.15	4676	500	500	no
WMAP3 J1601+3329	—	4C33.38	16 02 07.2	+33 26 53.0	Blazar Unknown-type/GPS	1.1	1656	900	900	no
WMAP3 J1608+1028	WMAP 9	4C10.45	16 08 46.0	+10 29 07.0	FSRQ	1.226	1412	2200	2200	yes
WMAP3 J1617−7716	WMAP 183	PKS1610-77	16 17 49.3	−77 17 17.9	FSRQ	1.71	3350	2000	1900	yes
WMAP3 J1613+3412	WMAP 23	OS319	16 13 40.9	+34 12 47.0	FSRQ	1.397	2324	3300	3300	yes
WMAP3 J1635+3807	WMAP 33	4C38.41	16 35 15.4	+38 08 03.9	FSRQ	1.813	3221	4900	4900	yes
WMAP3 J1637+4714	—	4C47.44	16 37 45.1	+47 17 32.9	FSRQ	0.74	1244	900	900	no
WMAP3 J1643−7712	—	PKS1637-77	16 44 16.0	−77 15 47.9	Radio Galaxy	0.043	2268	800	800	no
WMAP3 J1638+5722	WMAP 56	S41637+574	16 38 13.3	+57 20 24.0	FSRQ	0.75	1750	1500	1500	yes
WMAP3 J1633+8227	WMAP 76	NGC6251	16 32 26.1	+82 32 20.0	Radio Galaxy	0.024	1460	1400	1400	yes
WMAP3 J1642+3948	WMAP 35	3C345	16 42 58.6	+39 48 37.0	FSRQ	0.593	8719	6000	6000	yes
WMAP3 J1642+6853	WMAP 69	4C69.21	16 42 07.8	+68 56 38.0	FSRQ	0.751	1527	900	900	no
WMAP3 J1648+4114	—	GB6J1648+4104	16 48 29.2	+41 04 05.0	SSRQ	0.851	197	1200	900	no
WMAP3 J1651+0457	WMAP 10	HerA	16 51 08.1	+04 59 32.9	Radio Galaxy	0.154	11376	1100	1100	yes
WMAP3 J1654+3939	WMAP 36	MKR501	16 53 52.2	+39 45 35.9	BL Lac	0.034	1375	0	0	no
WMAP3 J1658+0742	WMAP 13	PKS1655+077	16 58 08.8	+07 41 27.9	FSRQ	0.621	1108	1000	1000	yes
WMAP3 J1656+5706	—	4C57.28	16 57 20.7	+57 05 53.0	FSRQ	1.281	764	600	600	no
WMAP3 J1657+4749	—	S41656+47	16 57 54.5	+47 49 14.9	FSRQ	0	0	600	600	no
WMAP3 J1703−6213	WMAP 198	PMNJ1703-6212	17 03 36.4	−62 12 39.9	FSRQ	0	616	2000	2000	no
WMAP3 J1659+6829	—	GB6J1700+6830	17 00 09.3	+68 30 06.0	FSRQ	0.301	380	700	700	no
WMAP3 J1707+0146	—	PKS1705+018	17 07 34.3	+01 48 45.0	FSRQ	2.568	463	900	900	no
WMAP3 J1716+6840	—	S41716+68	17 16 13.9	+68 36 38.0	FSRQ	0.777	838	700	700	no

Table 3: WMAP3 catalogue: basic data, identifications and inclusion in the flux limited sample (continued)

Source name IAU Convention	WMAP ID 1year catalogue	Common name	Ra(J2000.0)	Dec(J2000.0)	Class	Redshift	Radio Flux density 5 GHz (mJy)	WMAP flux density 41 GHz (mJy)	Corrected 41 GHz (mJy)	In Complete Sample
WMAP3 J1724−6500	WMAP 196	NGC6328	17 23 41.0	−65 00 36.0	Radio Galaxy/GPS	0.014	4373	1700	1700	yes
WMAP3 J1737−7934	WMAP 186	PKS1725-795	17 33 40.6	−79 35 54.9	Blazar Unknown-type	0	419	1100	1100	yes
WMAP3 J1727+4530	WMAP 43	S41726+45	17 27 27.6	+45 30 38.9	FSRQ	0.717	935	1100	1100	yes
WMAP3 J1734+3856	WMAP 38	1Jy1732+389	17 34 20.5	+38 57 51.0	FSRQ	0.97	557	1200	1200	yes
WMAP3 J1740+4739	—	S41738+476	17 39 57.1	+47 37 58.0	BL Lac	0	818	1100	1100	yes
WMAP3 J1740+5212	WMAP 48	4C51.37	17 40 36.9	+52 11 43.0	FSRQ	1.375	1699	1300	1300	yes
WMAP3 J1748+7006	WMAP 68	1Jy1749+701	17 48 32.8	+70 05 50.9	BL Lac	0.77	715	900	900	no
WMAP3 J1753+2848	WMAP 22	GB6J1753+2847	17 53 42.4	+28 48 03.9	Blazar Unknown-type	0	511	2400	2400	yes
WMAP3 J1753+4405	—	S41751+441	17 53 22.6	+44 09 46.0	FSRQ	0.871	1000	800	800	no
WMAP3 J1803−6507	WMAP 199	PKS1758-651	18 03 23.5	−65 07 36.0	Blazar Unknown-type	0	780	1500	1500	yes
WMAP3 J1758+6632	WMAP 64	NGC6543	17 58 33.4	+66 37 58.0	Planetary Nebula	0	875	600	600	no
WMAP3 J1759+3852	—	S41758+3	18 00 24.7	+38 48 29.9	FSRQ	2.092	735	800	800	no
WMAP3 J1801+4404	—	1Jy1800+440	18 01 32.2	+44 04 22.0	FSRQ	0.663	1193	1400	1400	yes
WMAP3 J1800+7828	WMAP 72	S51803+784	18 00 45.7	+78 28 04.0	BL Lac	0.68	2620	1700	1700	yes
WMAP3 J1806+6949	WMAP 67	3C371	18 06 50.3	+69 49 27.9	BL Lac	0.051	2122	1200	1200	yes
WMAP3 J1820−6343	WMAP 200	PKS1814-63	18 19 34.9	−63 45 47.9	Radio Galaxy	0.063	4506	1200	1200	yes
WMAP3 J1820−5521	—	PKS1815-553	18 19 45.4	−55 21 20.0	FSRQ	0	1124	0	0	no
WMAP3 J1824+5650	WMAP 53	4C56.27	18 24 07.0	+56 51 01.0	BL Lac	0.664	1263	1400	1400	yes
WMAP3 J1829+4844	WMAP 46	3C380.0	18 29 31.7	+48 44 45.9	SSRQ	0.692	5519	2500	2500	yes
WMAP3 J1837−7106	WMAP 192	PKS1831-711	18 37 28.6	−71 08 43.0	FSRQ/GPS	1.356	2294	1500	1500	yes
WMAP3 J1835+3246	—	3C382.0	18 35 02.1	+32 41 49.9	Radio Galaxy	0.058	2281	700	700	no
WMAP3 J1842+6807	WMAP 66	S41842+68	18 42 33.6	+68 09 24.0	FSRQ	0.472	925	1200	1200	yes
WMAP3 J1840+7946	WMAP 73	3C390.3	18 42 09.0	+79 46 17.0	SSRQ	0.056	4380	1000	1000	yes
WMAP3 J1848+3221	—	B21846+32A	18 48 22.0	+32 19 01.9	FSRQ	0.798	762	800	800	no
WMAP3 J1850+2823	WMAP 28	S21848+28	18 50 27.4	+28 25 12.0	FSRQ/GPS	2.56	1013	900	900	no
WMAP3 J1849+6705	WMAP 65	4C66.20	18 49 15.9	+67 05 39.9	FSRQ	0.657	845	1400	1400	yes
WMAP3 J1902+3153	WMAP 34	3C395	19 02 55.9	+31 59 42.0	FSRQ	0.635	1863	800	800	no
WMAP3 J1923−2106	WMAP 8	PMNJ1923-2104	19 23 32.2	−21 04 32.9	FSRQ	0.874	2885	2100	2100	yes
WMAP3 J0000+0000	—	OV-236	19 24 51.4	−29 14 30.0	FSRQ	0.352	14332	13200	13200	yes
WMAP3 J1927+6118	WMAP 59	S41926+611	19 27 30.4	+61 17 31.9	BL Lac	0	558	1100	1100	yes
WMAP3 J1927+7357	WMAP 70	4C73.18	19 27 48.4	+73 58 00.9	FSRQ	0.302	3626	2900	2900	yes
WMAP3 J1937−3956	—	PKS1933-400	19 37 16.3	−39 58 00.9	FSRQ	0.965	1129	1100	1100	yes
WMAP3 J1938−6345	—	PKS1934-63	19 39 25.0	−63 42 44.9	Blazar Unknown-type/GPS	0.183	5910	0	0	no
WMAP3 J1952+0233	—	3C403.0	19 52 15.7	+02 30 24.0	Radio Galaxy	0.059	2026	600	600	no
WMAP3 J1955+5132	WMAP 51	S41954+513	19 55 42.7	+51 31 49.0	FSRQ	1.22	1675	800	800	no
WMAP3 J1958−3845	WMAP 3	PKS1954-388	19 57 59.8	−38 45 06.0	FSRQ	0.63	1944	3000	3000	yes
WMAP3 J2000−1749	WMAP 11	PKS1958-179	20 00 57.1	−17 48 56.9	FSRQ	0.65	2700	1600	1600	yes
WMAP3 J2005+7755	—	S52007+777	20 05 31.1	+77 52 42.9	BL Lac	0.342	1260	0	0	no
WMAP3 J2011−1547	WMAP 14	PKS2008-159	20 11 15.7	−15 46 40.0	FSRQ	1.18	949	1500	1500	yes
WMAP3 J2010+7230	—	4C72.28	20 09 52.4	+72 29 18.9	BL Lac	0	907	800	800	no
WMAP3 J2022+6136	WMAP 63	S42021+61	20 22 06.7	+61 36 57.9	Radio Galaxy	0.227	2743	1300	1300	no
WMAP3 J2035−6845	WMAP 194	PKS2030-689	20 35 48.7	−68 46 32.9	FSRQ	1.084	648	800	800	no
WMAP3 J2035+1058	—	PKS2032+107	20 35 22.3	+10 56 07.0	BL Lac	0.601	759	0	0	no
WMAP3 J2056−4717	WMAP 208	PKS2052-47	20 56 16.3	−47 14 47.0	FSRQ	1.489	2026	1900	1900	yes
WMAP3 J2109−4114	WMAP 1	PKS2106-413	21 09 33.1	−41 10 19.9	FSRQ	1.058	2076	1200	1200	yes
WMAP3 J2109+3537	WMAP 49	GB6J2109+3532	21 09 31.9	+35 32 57.0	Unidentified	0	1208	700	700	no
WMAP3 J2123+0536	WMAP 27	PKS2121+053	21 23 44.5	+05 35 21.9	FSRQ	1.878	1930	1700	1700	yes
WMAP3 J2131−1206	WMAP 17	PKS2128-12	21 31 35.2	−12 07 05.9	FSRQ	0.501	2989	2300	2300	yes
WMAP3 J2134−0154	WMAP 20	4C-02.81	21 34 10.3	−01 53 17.0	BL Lac	1.285	1308	1700	1700	yes
WMAP3 J2136+0042	WMAP 25	PKS2134+004	21 36 38.5	+00 41 53.9	FSRQ	1.932	8416	3000	3000	yes
WMAP3 J2139+1424	WMAP 41	PKS2136+141	21 39 01.3	+14 23 35.0	FSRQ	2.427	1073	1900	1900	yes
WMAP3 J2148−7757	WMAP 184	PKS2141-781	21 46 29.9	−77 55 54.0	FSRQ	0.334	508	0	0	no
WMAP3 J2143+1740	WMAP 44	S32141+17	21 43 35.5	+17 43 49.0	FSRQ	0.211	1006	0	0	no
WMAP3 J2148+0657	WMAP 37	4C06.69	21 48 05.3	+06 57 38.9	FSRQ	0.99	4949	7200	7200	yes
WMAP3 J2151−3026	—	PKS2149-307	21 51 55.4	−30 27 52.9	FSRQ	2.345	1336	1200	1200	yes
WMAP3 J2157−6942	WMAP 190	PKS2153-69	21 57 05.9	−69 41 23.9	SSRQ	0.028	11649	2600	2600	yes
WMAP3 J2158−1502	WMAP 18	PKS2155-152	21 58 06.1	−15 01 09.0	FSRQ	0.672	3091	1700	1700	yes
WMAP3 J2202+4217	WMAP 58	BLLAC	22 02 43.2	+42 16 40.0	BL Lac	0.069	2940	3100	3100	no

Giommi et al.: A WMAP complete sample of blazars

Table 3: WMAP3 catalogue: basic data, identifications and inclusion in the flux limited sample (continued)

Giommi et al.: A WMAP complete sample of blazars

Source name IAU Convention	WMAP ID 1year catalogue	Common name	Ra(J2000.0)	Dec(J2000.0)	Class	Redshift	Radio Flux density 5 GHz (mJy)	WMAP flux density 41 GHz (mJy)	Corrected 41 GHz (mJy)	In Complete Sample
WMAP3 J2203+3146	WMAP 54	4C31.63	22 03 15.0	+31 45 38.0	FSRQ	0.295	2806	2200	2200	yes
WMAP3 J2203+1724	WMAP 45	PKS2201+171	22 03 26.8	+17 25 47.9	FSRQ	1.075	834	1700	1700	yes
WMAP3 J2206-1839	WMAP 16	PKS2203-18	22 06 10.3	-18 35 39.0	FSRQ	0.619	4271	1300	1300	yes
WMAP3 J2207-5349	—	PKS2204-54	22 07 43.6	-53 46 33.9	FSRQ	1.206	1410	800	800	no
WMAP3 J2211+2351	WMAP 50	PKS2209+236	22 12 05.9	+23 55 40.0	FSRQ/GPS	1.125	1123	1400	1400	yes
WMAP3 J2218-0335	WMAP 30	4C-03.79	22 18 52.0	-03 35 36.9	FSRQ	0.901	2717	2000	2000	yes
WMAP3 J2225-0456	WMAP 29	3C446	22 25 47.2	-04 57 01.0	FSRQ	1.404	6382	4100	4100	yes
WMAP3 J2229-0832	WMAP 24	PKS2227-08	22 29 40.0	-08 32 53.9	FSRQ	1.559	2423	2600	2600	yes
WMAP3 J2229-2051	—	PMNJ2229-2049	22 29 48.1	-20 49 26.0	Planetary Nebula	0	326	800	800	no
WMAP3 J2232+1144	WMAP 47	4C-11.69	22 32 36.4	+11 43 50.0	FSRQ	1.037	3967	3400	3400	yes
WMAP3 J2235-4834	WMAP 206	PKS2232-488	22 35 13.1	-48 35 57.9	FSRQ	0.51	1104	1800	1800	yes
WMAP3 J2236+2825	WMAP 57	B22234+28A	22 36 22.3	+28 28 58.0	FSRQ	0.795	1625	1100	1100	yes
WMAP3 J2239-5700	WMAP 201	PKS2236-572	22 39 12.1	-57 01 00.0	Unidentified	0	1063	0	0	no
WMAP3 J2246-1210	WMAP 21	PKS2243-123	22 46 18.1	-12 06 51.9	FSRQ	0.632	2661	1100	1100	yes
WMAP3 J2254+1608	WMAP 55	3C454.3	22 53 57.7	+16 08 53.0	FSRQ	0.859	14468	6800	6800	yes
WMAP3 J2255+4202	—	S42253+41	22 55 36.7	+42 02 50.9	FSRQ	1.476	1084	900	800	no
WMAP3 J2256-2011	WMAP 19	PKS2254-204	22 56 41.0	-20 11 40.9	BL Lac	0	454	800	800	no
WMAP3 J2258-2757	WMAP 12	PKS2255-282	22 58 05.8	-27 58 18.9	FSRQ	0.926	2127	6600	6600	yes
WMAP3 J2315-5017	WMAP 204	PKS2312-505	23 15 44.2	-50 18 38.9	Unidentified	0	341	900	900	no
WMAP3 J2322+5107	—	GB6J2322+5057	23 22 25.9	+50 57 51.0	FSRQ	1.279	1305	800	800	no
WMAP3 J2327+0939	—	PKS2325+093	23 27 33.4	+09 40 09.0	FSRQ	1.843	886	1200	1200	yes
WMAP3 J2329-4732	—	PKS2326-477	23 29 17.7	-47 30 19.0	FSRQ	1.299	2206	1400	1400	yes
WMAP3 J2330+1056	—	4C10.73	23 30 40.7	+11 00 18.0	FSRQ	1.489	1194	1000	1000	yes
WMAP3 J2331-1559	WMAP 32	PKS2329-16	23 31 38.5	-15 56 57.0	FSRQ	1.153	1879	800	800	no
WMAP3 J2333-2340	—	PKS2331-240	23 33 55.2	-23 43 40.0	Radio Galaxy	0.048	907	1200	1200	yes
WMAP3 J2334+0734	—	GB6J2334+0736	23 34 12.7	+07 36 25.9	FSRQ	0.401	953	1100	1100	yes
WMAP3 J2335-0129	—	PKS2332-017	23 35 20.3	-01 31 09.0	FSRQ	1.184	581	1300	1300	yes
WMAP3 J2335-5242	WMAP 195	PKS2333-528	23 36 12.1	-52 36 21.9	FSRQ	0	1588	500	400	no
WMAP3 J2346+0927	—	4C09.74	23 46 36.7	+09 30 44.9	FSRQ	0.677	1344	700	600	no
WMAP3 J2348-1631	WMAP 39	PKS2345-16	23 48 02.5	-16 31 11.9	FSRQ	0.6	2360	2000	2000	yes
WMAP3 J2354+4550	WMAP 74	4C45.51	23 54 21.7	+45 53 03.9	FSRQ	1.992	1127	1300	1300	yes
WMAP3 J2356+4952	WMAP 75	1Jy2352+495	23 55 09.4	+49 50 08.0	Radio Galaxy	0.238	2306	0	0	no
WMAP3 J2357-5313	WMAP 189	PKS2355-534	23 57 53.2	-53 11 12.9	FSRQ	1.006	1782	1100	1100	yes
WMAP3 J2358-1013	—	PKS2355-106	23 58 10.8	-10 20 08.0	FSRQ	1.639	1618	0	0	no
WMAP3 J2358-6050	WMAP 187	PKS2356-61	23 58 45.5	-60 52 54.9	Radio Galaxy	0.096	4421	1200	1200	yes